

Rheology Cyberinfrastructure for Integrated Research and Learning at ARCo7

TECHNICAL UNIVERSITY BERLIN, GERMANY
MARCH 20-21, 2007

The fourth Amherst Rheology Course (ARCo7) met in Berlin/Germany at the Institute of Manfred Wagner at the Technical University Berlin. The theme of the course was “synergy between experiments and theory in rheology”. ARCo7 provided participants with a broad overview of experimental and theoretical rheology and then equipped them with the cyberinfrastructure (CI) tools to obtain quantitative answers in rheology. Lectures and practice sessions were combined to develop a deeper understanding for the underlying

ing concepts that lead to quantitative results.

Rheology is an essential tool for solving complex problems at the frontiers of materials science. To be effective, materials scientists and engineers should be able to easily exchange rheological data and to merge data with theoretical viewpoints. We anticipate that, in the near future, companies will take advantage of CI and apply rheology more widely and efficiently. The company with the most effective CI tools and best CI-trained work force will be the most competitive.

© Appl. Rheol. 17 (2007) 302–304

The CI tools of ARCo7 consist of a single workspace (“rheology hub”) that bundles expert codes on the PC. This system allows the world’s leading rheology experts to easily share their expertise with a wide community of users. Implementation of such single workspace technology will introduce a new style of networking between individual researchers and/or between company sites. It allows for screening of materials, optimizing of processes, and evaluating hypothetical materials before even making them. Researchers and practitioners will routinely merge experimental data with modeling predictions and simulation (laboratory internal modeling tools, third party codes, free software). This merger will be further supported by the unified data standard that facilitates access anywhere-anytime. Global teleconferences and sales negotiations will be strengthened with rapid retrieval of material data and with ‘on the spot’ modeling calculations. Quantitative technical information will be retrieved or quickly generated (by modeling) for rapid decision-making. Practitioners of rheology in industry will evaluate their strengths locally by exploring their own ideas and hypotheses in the context of global rheology knowledge. Implementation of most of this CI technology is possible with today’s tools [1].

This spring, ARCo7 participants brought their own laptop computers and performed practice calculations with CI tools. Course participants used the “rheology hub” to compare experiments (dynamic mechanical, steady shear, startup of shear, startup various extensional flows, molecular weight distribution) with predictions from a range of theories, including classical theories (Maxwell, Rouse, Lodge, Doi-Edwards) and three recent polymer dynamics theories: the ‘tube dilation’ model of McLeish and coworkers [2 - 7], the hierarchical model of Larson and coworkers [8, 9] and the “molecular stress function” model of Wagner and coworkers [10]. The “NAPLES” code of Masubuchi and coworkers [11 - 13] was used to simulate molecular dynamics of homogeneous mixtures of molecules with different architecture. Furthermore, we practiced the new module for predicting the (monomodal, bimodal) molecular weight distribution of linear polymers from their dynamic mechanical data [14, 15]. Time-resolved rheometry tools [16] were applied to chemical gelation. Dynamic mechanical data were shifted into master curves and the (discrete and continuous) relaxation time spectra were calculated [17,

18]. In addition, Jonathan Rothstein expanded the scope by introducing into “Micro-Rheology” and “Rheology of Micellar Fluids”. At the end of the 2-day course, participants were able to intuitively and seamlessly navigate the “rheology hub”. Even course participants arriving at the course with minimal background in rheology efficiently navigated the rheology workspace.

The current group of rheology users worldwide is still fairly small and specialized, while the main workforce in materials science and engineering has little access to rheology. The Amherst Rheology Courses seek to broaden and diversify the population of individuals and institutions participating in rheological activities and to integrate rheology into wider areas of science and engineering. One reason for rheology’s current insular state is the lack of intuitive tools for analyzing rheological experiments and for comparing experiments with theory. A second reason is the lack of a universally accepted data standard. These problems are being addressed by using the “rheology hub” and by teaching its use to a wider group of people.

Twenty-one rheologists participated in ARCo7. The course was jointly directed and taught by H. Henning Winter (USA) and Manfred Wagner. Jonathan Rothstein (USA) joined the team as one of the instructors. The course started on March 19, 2007, with a welcome reception where the course participants got acquainted with one another. It ended with joint exercises and discussions on March 21. The Amherst Rheology Course (ARCo8) will convene again in June 2008, in Amherst, Massachusetts, USA. New topics will be added concerning the rheology of emulsions. Information about the “rheology hub” and about future ARC can be found at <http://rheology.tripod.com/ARC.htm>.

REFERENCES

- [1] Winter HH, Mours M: The cyber infrastructure initiative for rheology. *Rheol. Acta* 45 (2006) 331-338.
- [2] McLeish TCB, Larson RG: Molecular constitutive equations for a class of branched polymers: the pom-pom polymer. *J Rheol.* 42 (1998) 81-110.
- [3] McLeish TCB, Allgaier J, Bick DK, Bishko G, Biswas P, Blackwell R, Blottiere B, Clarke N, Gibbs B, Groves DJ, Hakiki A, Hoenan RK, Johnson JM, Kant R, Read DJ, Young RN: Dynamics of entangled H-polymers: theory, rheology, and neutron scattering. *Macromolecules* 32 (1999) 6734-6758.
- [4] Milner ST, McLeish TCB: Parameter-free theory for stress relaxation in star polymer melts.

- Macromolecules 30 (1997) 2159-2166.
- [5] Milner ST; McLeish, TCB: Reptation and contour-length fluctuations in melts of linear polymers. Phys. Rev. Lett. 81 (1998) 725-728.
 - [6] Blackwell RJ, Harlen OG, McLeish TCB: Theoretical linear and nonlinear rheology of symmetric tree-like polymer melts. Macromolecules 34 (2001) 2579-2596.
 - [7] Pryke A, Blackwell R J, McLeish TCB, Young RN: Synthesis, hydrogenation, and rheology of 1,2-polybutadiene star polymers. Macromolecules 35 (2002) 467-472.
 - [8] Larson RG: Combinatorial rheology of branched polymer melts. Macromolecules 34 (2001) 4556.
 - [9] Park SJ, Shanbhag S, Larson RG: A hierarchical algorithm for predicting the linear viscoelastic properties of polymer melts with long-chain branching. Rheol. Acta 44 (2005) 319-330.
 - [10] Wagner MH, Yamaguchi M, Takahashi M: Quantitative assessment of strain hardening of low-density polyethylene melts by the molecular stress function model. J. Rheol. 47 (2003) 779-793.
 - [11] Masubuchi Y, Ianniruberto G, Greco F, Marrucci G: Entanglement molecular weight and frequency response of sliplink networks. J. Chem. Phys. 119 (2003) 6925-6930.
 - [12] Masubuchi Y, Ianniruberto G, Greco F, Marrucci G: Molecular simulations of longtime behavior of entangled polymeric liquids by the primitive chain network model, modelling simulation. Mat. Sci. Eng. 12 (2004) 91-100.
 - [13] Masubuchi Y, Takimoto J, Koyama K, Ianniruberto G, Greco F, Marrucci G: Brownian Simulations of a Network of Reptating Primitive Chains. J. Chem. Phys. 115 (2001) 4387-4394.
 - [14] Nobile MR, Cocchini F: Evaluation of molecular weight distribution from dynamic moduli. Rheol. Acta 40 (2001) 111-119.
 - [15] Cocchini F, Nobile MR: Constrained inversion of rheological data to molecular weight distribution for polymer melts. Rheol. Acta 42 (2003) 232-242.
 - [16] Mours M, Winter HH: Time resolved rheometry. Rheol. Acta 33 (1994) 385-397.
 - [17] Baumgärtel M, Winter HH: Interrelation between continuous and discrete relaxation time spectra. J Non-Newtonian Fluid Mech 44 (1992) 15-36.
 - [18] Jackson JK, De Rosa ME, Winter HH: Molecular weight dependence of relaxation time spectra for the entanglement and flow behavior of monodisperse linear flexible polymers. Macromolecules 27 (1994) 2426-2431.


H. Henning Winter
Amherst MA, USA
winter@ecs.umass.edu



IRIS drives innovation in rheology: interactive graphics for data analysis, seamless communication of data, comparison with models and more. Import data from any source, shift, calculate spectra, compare, store, retrieve, and plot within minutes.

IRIS allows the rheologist to pursue his/her real mission: explore new materials, discover relaxation patterns, apply to processes, be quantitative and reliable, communicate results, explain and teach.

IRIS (new) predicts from molecular theory: (a) tube dilation theory of T. McLeish and coworkers and (b) molecular stress function theory of M. Wagner.



E-mail: IRISrheo@yahoo.com
<http://rheology.tripod.com/>