

# PHYSICAL AND BIOLOGICAL STUDY OF CELL CULTURES IN A BIOREACTOR: ON-LINE AND OFF-LINE RHEOLOGICAL ANALYSES

YANNICK MANON<sup>1</sup>, DOMINIQUE ANNE-ARCHARD<sup>3</sup>, JEAN-LOUIS URIBELARREA<sup>1</sup>, CAROLE MOLINA-JOUE<sup>1</sup> AND LUC FILLAUDEAU<sup>1-3\*</sup>

<sup>1</sup> Université de Toulouse, INRA & CNRS & INSA, LISBP (Laboratoire d'Ingénierie des Systèmes Biologiques et des Procédés), 135 avenue de Rangueil, 31077 Toulouse, France

<sup>2</sup> Université de Toulouse, INPT-UPS & CNRS, IMFT (Institut de Mécanique des Fluides de Toulouse), Allée Camille Soula, 31400 Toulouse, France

<sup>3</sup> CNRS, Fédération de Recherche FERMAT, 31400 Toulouse, France

\* Corresponding author: [luc.fillaudeau@insa-toulouse.fr](mailto:luc.fillaudeau@insa-toulouse.fr)  
Fax: x33.5.61559400

Received: 7.12.2010, Final version: 19.2.2011

## ABSTRACT:

Rheological behaviour of culture broth stands as a fundamental parameter in bioprocess performances because it affects simultaneously the heat and mass transfer as well as the flow pattern. On-line measurements of rheological behaviour are hardly compatible with the operating condition with respect to accurate and stringent conditions imposed by cell culture strategy. Our scientific and technical objectives are (i) to develop and identify an experimental device enabling on-line rheometry and (ii) to discuss and compare on-line and off-line measurements. In this aim, a bioreactor was equipped with a derivation loop including a specific on-line rheometric device as well as additional physical and biological measurements (specific density, mass flow rate, electrical conductivity, pH, pO<sub>2</sub> and temperature) during microbial cell cultures. In a first time, friction curves of calibrated ducts were established with Newtonian and non-Newtonian shear-thinning fluids. In a second time, axenic cultures with two microorganisms (bacteria and yeast exhibiting different sizes) were investigated in pure oxidative culture in order to produce biomass under high cell concentrations: ~ 40 to 110 g<sub>CDW</sub>/l for *E. coli* (bacteria) and ~ 75 to 105 g<sub>CDW</sub>/l for *Y. lipolytica* (yeast). Cell broths exhibited Newtonian behaviour for *E. coli* and shear-thinning behaviour for *Y. lipolytica*, which were both dependant on biomass concentration. On-line and off-line rheological measurements are consistent for *E. coli* and *Y. lipolytica*, but significantly differed. On-line estimated viscosity appears higher than off-line apparent viscosity. Several assumptions in relation with microorganism physiology and metabolism (size, morphology, surface properties, concentration, biological activity) could be formulated in agreement with scientific literature. On-line rheology brings new insight to investigate complex interaction between physical and biological phenomena.

## ZUSAMMENFASSUNG:

Das rheologische Verhalten von Kulturen stellt einen entscheidenden Parameter in der Leistungsfähigkeit von biologischen Prozessen dar, da es sowohl die Wärme und Massenübertragung, als auch das Flussverhalten beeinflusst. On-line Messungen des rheologischen Verhaltens sind nicht mit den exakten Arbeitsbedingungen die für die Zellkulturstategie nötig sind, vereinbar. Unsere wissenschaftlichen und technischen Ziele sind (i) die Entwicklung eines experimentellen Instruments zur on-line Rheometrie und (ii) on-line und off-line Messungen miteinander zu vergleichen. Dafür wurde ein Bioreaktor mit einem Rohr in Derivation mit einem spezifischen on-line Rheometrie Instrument und zusätzlichen physikalischen und biologischen Meßgeräten (spezifische Dichte, Durchflussrate, elektrische Leitfähigkeit, pH, pO<sub>2</sub> und Temperatur) ausgestattet. Zunächst wurden Reibwertkurven erstellt mit kalibrierten Mitteln mit Newtonsches und allgemeinen Nicht-Newtonsches strukturviskosen Flüssigkeiten. Im zweiten Schritt wurden axenische Kulturen von zwei Mikroorganismen (Bakterien und Hefen verschiedener Größen) in reiner oxidativen Kultur untersucht um Biomasse mit hoher Zellkonzentration zu erzeugen: ~ 40–110 g<sub>CDW</sub>/l für *E. coli* (Bakterien) und ~ 75–105 g<sub>CDW</sub>/l für *Y. lipolytica* (Hefe). Das Kulturmedium zeigte Newtonsches Verhalten mit *E. coli* und strukturviskoses Verhalten bei *Y. lipolytica*, wobei beides von der Biomassekonzentration anhing. On-line und off-line rheologische Messungen sind consistent für *E. coli* und *Y. lipolytica*, aber unterscheiden sich deutlich. Die geschätzten Werte für Viskosität sind höher wenn on-line gemessen. Verschiedene Annahmen über die Physiologie des Mikroorganismus und den Metabolismus (Größe, Morphologie, Oberflächeneigenschaften, Konzentration, biologische Aktivität) konnten im Einklang mit der Literatur formuliert werden. On-line Rheologie ermöglicht neue Einblicke in die Untersuchung der komplexen Interaktion zwischen physikalischen und biologischen Phänomenen.

## RÉSUMÉ:

Le comportement rhéologique des mouts de fermentation représente un paramètre fondamental pour le contrôle et l'intensification des bioprocédés car il affecte simultanément les transferts de chaleur, de matière et de

© Appl. Rheol. 21 (2011) 35167

DOI: 10.3933/AppRheol-21-35167

This is an extract of the complete reprint-pdf, available at the Applied Rheology website

<http://www.appliedrheology.org>

35167-1

Applied Rheology  
Volume 21 · Issue 3

This is an extract of the complete reprint-pdf, available at the Applied Rheology website

<http://www.appliedrheology.org>

behaviour [8] and it may not be excluded that these wall properties evolved when the cell culture conditions are not respected, as occurs when a sample is taken. Supernatant properties can also affect the broth behaviour when cells produce metabolites or extracellular polymeric substances [17]. But this third assumption is not true here as the supernatant of broth cultures is Newtonian with very low viscosity. However, all these works are usually realised off-line and do not take into account the impact of biological activity and the presence of gas volume fraction (aeration/respiration) in broth. This point needs to be taken into account as shown by Barbot et al. [28] who explored the influence of respiration in the case of activated sludge under aerated condition within a bioreactor. In their experiment, rheological behaviour was evaluated using on-line torque measurements. They observed that inhibition of respiration by sodium azide addition leads to an evolution of the rheological behaviour. However, the comparison between activated sludge and axenic cell cultures is limited considering differences between experimental conditions and broth composition. But this demonstrates that biological activity is an additional difficulty for broth characterization and has to be considered during the measurements.

#### 4 CONCLUSION

Rheological behaviour of broth stands as a fundamental parameter that could limit bioprocess performances. Therefore axenic culture of microorganism in bioreactor involves culture strategy with aeration as a necessary and obligatory parameter. All the complexity related to on-line monitoring of rheological behaviour results in the setting up of an adapted tool with correct measurement in respect of cell culture conditions. In the case of torque measurement, the variation of agitation speed and aeration flow rate to allow correct flow regime can hardly be achieved (risk of disrupt the system) except in robust bioprocess (wastewater treatment).

In this study, our objectives were to investigate on-line and off-line physical measurement with a new device during axenic cell cultures under high biomass concentration. This dedicated and innovative experimental set-up was developed and was calibrated with well-defined Newtonian and non-Newtonian fluids. Two

microorganisms (*E. coli* and *Y. lipolytica*) with different cell size were cultivated. The physical measurements showed different results: (i) the impact of cell concentration on pressure drop, (ii) the contribution of the gas phase in the broth, (iii) the possible impact of difference between off-line and on-line shear rate ranges that can be explored and (iv) a setting up of caution to use properly our experimental set-up and to estimate the on-line viscosity equivalent to a homogenous liquid medium. Our works demonstrated the need and usefulness of on-line rheology measurements to achieve and quantify reliable information about the dynamic rheological behaviour of cell broths during cultures. Our researches bring new insights on the dynamical interactions between physical and biological phenomena in bioprocess which will be useful to progress in the evaluation and the control of their performances.

#### ACKNOWLEDGMENTS

Authors are grateful to Fédération de recherche FERMaT FR3089 INPT-UT3-CNRS-INSA TOULOUSE and Bérange PORTELLI (Ph. D. Student, LISBP) for scientific and technical contribution.

#### REFERENCES

- [1] Cascaval D, Oniscu C, Galaction AI: Rheology of fermentation broths. 2 – Influence of the rheological behavior on biotechnological processes, Rev. Roum. Chim. 48 (2003) 339–356.
- [2] Pamboukian CRD, Facciotti MCR: Rheological and morphological characterization of *Streptomyces olindensis* growing in batch and fed-batch fermentations, Braz. J. Chem. Eng. 22 (2005) 31–40.
- [3] Petersen N, Stocks S, Gernaey KV: Multivariate models for prediction of rheological characteristics of filamentous fermentation broth from the size distribution, Biotechnol. Bioeng. 100 (2008) 61–71.
- [4] Corrieu G: The influence of rheological behavior on the performance of bioreactors, Biofutur (1997) A2–A4.
- [5] Ballica R, Ryu DDY, Powell RL, Owen D: Rheological Properties Of Plant-Cell Suspensions, Biotechnol. Prog. 8 (1992) 413–420.
- [6] Curtis WR, Emery AH: Plant-Cell Suspension-Culture Rheology, Biotechnol. Bioeng. 42 (1993) 520–526.
- [7] Raposo S, Lima-Costa ME: Rheology and shear stress of *Centaurea calcitrapa* cell suspension cultures grown in bioreactor, Biotechnol. Lett. 28 (2006) 431–438.
- [8] Jordan A, Duperray A, Verdier C: Fractal approach

- to the rheology of concentrated cell suspensions, *Phys. Rev. E* 77 (2008) 011911.
- [9] Trepap X, Deng LH, An SS, Navajas D, Tschumperlin DJ, Gerthoffer WT, Butler JP, Fredberg JJ: Universal physical responses to stretch in the living cell, *Nature* 447 (2007) 592–596.
- [10] Chen F, Chen H, Gong XD: Mixotrophic and heterotrophic growth of *Haematococcus lacustris* and rheological behaviour of the cell suspensions, *Bioresour. Technol.* 62(1997) 19–24.
- [11] Cho YJ, Hwang HJ, Kim SW, Song CH, Yun JW: Effect of carbon source and aeration rate on broth rheology and fungal morphology during red pigment production by *Paecilomyces sinclairii* in a batch bioreactor, *J. Biotechnol.* 95 (2002) 13–23.
- [12] Rodríguez Porcel EM, Casas López JL, Sánchez Pérez JA, Fernández Sevilla JM, Chisti Y: Effects of pellet morphology on broth rheology in fermentations of *Aspergillus terreus*, *Biochem. Eng. J.* 26 (2005) 139–144.
- [13] Gogus N, Tari C, Oncu S, Unluturk S, Tokatli F: Relationship between morphology, rheology and polygalacturonase production by *Aspergillus sojae* ATCC 20235 in submerged cultures, *Biochem. Eng. J.* 32 (2006) 171–178.
- [14] Mancini M, Moresi M: Rheological behaviour of baker's yeast suspensions, *J. Food Eng.* 44 (2000) 225–231.
- [15] Klein J, Maia J, Vicente AA, Domingues L, Teixeira JA, Jurascik M: Relationships between hydrodynamics and rheology of flocculating yeast suspensions in a high-cell-density airlift bioreactor, *Biotechnol. Bioeng.* 89 (2005) 393–399.
- [16] Toda K, Furuse H, Amari T, Wei XF: Cell concentration dependence of dynamic viscoelasticity of *Escherichia coli* culture suspensions, *J. Ferment. Bioeng.* 85 (1998) 410–415.
- [17] O'Cleirigh C, Casey JT, Walsh PK, O'Shea DG: Morphological engineering of *Streptomyces hygroscopicus* var. *geldanus*: regulation of pellet morphology through manipulation of broth viscosity, *Appl. Microbiol. Biotechnol.* 68 (2005) 305–310.
- [18] Borges CD, Vendruscolo CT: Xanthan synthesized by strains of *Xanthomonas campestris* pv *pruni*: production, viscosity and chemical composition, *Bioscience Journal* 23 (2007) 67–73.
- [19] Porcel EMR, Lopez JLC, Perez JAS, Sevilla JMF, Chisti Y: Effects of pellet morphology on broth rheology in fermentations of *Aspergillus terreus*, *Biochem. Eng. J.* 26 (2005) 139–144.
- [20] Al-Asheh S, Abu-Jdayil B, Abunasser N, Barakat A: Rheological characteristics of microbial suspensions of *Pseudomonas aeruginosa* and *Bacillus cereus*, *Int. J. Biol. Macromol.* 30 (2002) 67–74.
- [21] Kembrowski ZBPOP: On-line measurements of the rheological properties of fermentation broth, *Rheol. Acta* 29 (1990) 588–593.
- [22] Seyssiecq I, Marrot B, Djerroud D, Roche N: In situ triphasic rheological characterisation of activated sludge, in an aerated bioreactor, *Chem. Eng. J.* 142 (2008) 40–47.
- [23] Shah RK, London AL: Laminar flow forced convection in ducts: a source book for compact heat exchanger analytical data, *Advances in Heat Transfer*, Academic Press, New York (1978).
- [24] Churchill SW: Friction-Factor Equation Spans All Fluid-Flow Regimes, *Chem. Eng.* 84 (1977) 91–92.
- [25] Metzner AB, Reed JC: Flow of non-newtonian fluids-correlation of the laminar, transition, and turbulent-flow regions, *Aiche J.* 1 (1955) 434–440.
- [26] OIML-GUM: Evaluation of measurement data – Guide to the expression of uncertainty in measurement, Joint Committee on Guides in Metrology OIML G 1-100 Edition 2008, Corrected version 2010.
- [27] Wood JM: Osmosensing by bacteria: Signals and membrane-based sensors, *Microbiology And Molecular Biology Reviews* 63 (1999) 230–262.
- [28] Barbot E, Seyssiecq I, Roche N, Marrot B: Inhibition of activated sludge respiration by sodium azide addition: Effect on rheology and oxygen transfer, *Chem. Eng. J.* 163 (2010) 230–235.

